arXiv:1511.03679v1 [math-ph] 11 Nov 2015

V.V. Borzov [1], **E.V. Damaskinsky** [2]

Invariance of the generalized oscillator under linear transformation of the related system of orthogonal polynomials

We consider two families of polynomials $\mathbb{P} = \{P_n(x)\}_{n=0}^{\infty}$ and $\mathbb{Q} = \{Q_n(x)\}_{n=0}^{\infty}$ [3] orthogonal on the real line with respect to probability measures μ and ν respectively. Let $\{Q_n(x)\}_{n=0}^{\infty}$ and $\{P_n(x)\}_{n=0}^{\infty}$ connected by the linear relations

$$Q_n(x) = P_n(x) + a_1 P_{n-1}(x) + ... + a_k P_{n-k}(x).$$

Let us denote $\mathfrak{A}_P$ and $\mathfrak{A}_Q$ generalized oscillator algebras associated with the sequences $\mathbb{P}$ and $\mathbb{Q}$. In the case $k = 2$ we describe all pairs ($\mathbb{P}$,$\mathbb{Q}$), for which the algebras $\mathfrak{A}_P$ and $\mathfrak{A}_Q$ are equal. In addition, we construct corresponding algebras of generalized oscillators for arbitrary $k \geq 1$.

1 Introduction

Let $\mathbb{P} = \{P_n(x)\}_{n=0}^{\infty}$ is a family of polynomials orthogonal on the real line with respect to the probability measure μ. Consider the sequence of polynomials $\mathbb{Q} = \{Q_n(x)\}_{n=0}^{\infty}$ such that

$$Q_n(x) = P_n(x) + a_1 P_{n-1}(x) + ... + a_k P_{n-k}(x), \quad n > k-1.$$

The family of orthogonal polynomials associated with such linear relation was discussed in several works (see e.g. [1] - [5]). In particular the necessary and sufficient conditions for the orthogonality of the sequence $\{Q_n(x)\}_{n=0}^{\infty}$ with respect to a positive Borel measure ν on the real line are given in the article [5].

It is known [6] that every sequence of polynomials $\{P_n(x)\}_{n=0}^{\infty}$ orthogonal with respect to positive Borel measures μ on the real line generates the generalized oscillator algebra $\mathfrak{A}_P$. In this work, we investigate the question under what conditions algebras $\mathfrak{A}_P$ and $\mathfrak{A}_Q$, generated by such linearly related polynomials, coincide

$$\mathfrak{A}_P = \mathfrak{A}_Q.$$

This problem was considered in [8] for the simplest case $k = 1$. In this paper we discuss the case $k = 2$.

Below we will need the following results ([5]-[7]). Let u is a linear functional on the linear space of polynomials with real coefficients. The polynomials $\{P_n(x)\}_{n=0}^{\infty}$ are called orthogonal with respect to u, if

$$\langle u, P_n P_m \rangle = 0, \, \forall n \neq m \quad \text{and} \quad \langle u, P_n^2 \rangle \neq 0, \, \forall n = 0, 1, \ldots.$$

[1] St.Petersburg State University of Telecommunications, 191065, Moika 61, St.Petersburg, Russia. E-mail: borzov.vadim@yandex.ru

[2] VI(IT) affilated to VA MTO, Russia, 191123, Zacharievskaya 22, St.Petersburg, Russia. E-mail: evd@pdmi.ras.ru

[3] Here and below we consider only monic polynomials.

Let $H = \{u_{i+j}\}_{i,j\geq 0}$ is the Hankel matrix associated with the functional u, where $u_k = \langle u, x^k\rangle, k \geq 0$. The linear functional u is called the quasi-definite (positive definite) functional, if the leading submatrices H_n of the matrix H are nonsingular (positive definite) for all $n \geq 0$.

Favard's theorem gives a description of quasi-definite (positively definite) linear functional in terms of the three-term recurrence relations that satisfied by the polynomials $\{P_n(x)\}_{n=0}^{\infty}$:

$$xP_n(x) = P_{n+1}(x) + \beta_n P_n(x) + \gamma_n P_{n-1}(x)$$
$$P_0(x) = 1, \qquad P_1(x) = x - \beta_0,$$

where $\gamma_n \neq 0$ (respectively $\gamma_n > 0$).

If u is a positive definite linear functional, then there exists the positive Borel measure μ (supported by an infinite subset of $\mathbb{R}$) such that

$$\langle u, q\rangle = \int_{\mathbb{R}} q\mathrm{d}\mu, \qquad \forall \mathrm{q} \in \mathbb{R}.$$

Bellow we will use the following theorem.

Theorem 1.1. [5] *Let $\{P_n(x)\}_{n=0}^{\infty}$ is the sequence of orthogonal polynomials with recurrence relations*

$$xP_n(x) = P_{n+1}(x) + \beta_n P_n(x) + \gamma_n P_{n-1}, \quad (\gamma_n \neq 0) \tag{1}$$
$$P_0(x) = 1, \quad P_1(x) = x - \beta_0.$$

Let a_1 and a_2 are real numbers such that $a_2 \neq 0$, and $Q_n(x)$ are polynomials defined by the relations

$$Q_n(x) = P_n(x) + a_1 P_{n-1}(x) + a_2 P_{n-2}, \quad n \geq 3. \tag{2}$$

Then the orthogonality of the sequence $\{Q_n(x)\}_{n=0}^{\infty}$ depends on the choice of the coefficients a_1 and a_2. More precisely, $\{Q_n(x)\}_{n=0}^{\infty}$ is a family of orthogonal polynomials with recurrence relations

$$xQ_n(x) = Q_{n+1}(x) + \widetilde{\beta}_n Q_n(x) + \widetilde{\gamma}_n Q_{n-1}, \quad n \geq 1, \quad (\gamma_n \neq 0) \tag{3}$$

if and only if $\gamma_3 + a_1(\beta_2 - \beta_3) \neq 0$ *and*

(i) if $a_1 = 0$, then for $n \geq 4$, $\beta_n = \beta_{n-2}$, $\gamma_n = \gamma_{n-2}$;

(ii) if $a_1 \neq 0$ and $a_1^2 = 4a_2$, then for $n \geq 2$

$$\beta_n = A + Bn + Cn^2, \quad \gamma_n = D + En + Fn^2,$$

with $a_1 C = 2F$, $a_1 B = 2E - 2F$, $(A, B, C, D, E, F \in \mathbb{R})$;

(iii) if $a_1 \neq 0$ and $a_1^2 > 4a_2$, then for $n \geq 2$

$$\beta_n = A + B\lambda^n + C\lambda^{-n}, \quad \gamma_n = D + E\lambda^n + F\lambda^{-n}$$

with $a_1 C = (1+\lambda)F$, $a_1 \lambda B = (1+\lambda)E$, $(A, B, C, D, E, F \in \mathbb{R})$, *where λ is the unique solution in $(-1, 1)$ of the equation*

$$a_1^2 \lambda = a_2(1+\lambda)^2;$$

(iv) if $a_1 \neq 0$ *and* $a_1^2 < 4a_2$*, and we let* $\lambda = e^{i\theta}$*,* $\theta \in (0, \pi)$ *be the unique solution of the equation* $a_1^2\lambda = a_2(1+\lambda)^2$*, then for* $n \geq 2$

$$\beta_n = A + Be^{in\theta} + \overline{B}e^{-in\theta}, \quad \gamma_n = D + Ee^{in\theta} + \overline{E}e^{-in\theta},$$

with $a_1\lambda B = (1+\lambda)E \quad (A, D \in \mathbb{R},\ B, F \in \mathbb{C})$.

Let us give the definition of the generalized oscillator connected with the family of orthogonal polynomials [6]. Let μ is a probability measure on $\mathbb{R}$ with finite moments of all orders

$$\mu_n = \int_{-\infty}^{+\infty} x^n \mathrm{d}\mu < \infty, \quad \mathrm{n} \geq 0.$$

These moments define uniquely two sequences of real numbers $\{a_n\}_{n=0}^{\infty}$, $\{b_n\}_{n=0}^{\infty}$ and the system of orthogonal polynomials $\{\Psi_n(x)\}_{n=0}^{\infty}$ which satisfy recurrence relations

$$x\Psi_n(x) = b_n\Psi_{n+1}(x) + a_n\Psi_n(x) + b_{n-1}\Psi_{n-1}(x), \tag{4}$$

for $n \geq 0$ and also initial conditions

$$\Psi_0(x) = 1, \qquad \Psi_1(x) = \frac{x - a_0}{b_0}.$$

These polynomials form an orthonormal basis in the Hilbert space $\mathcal{H} = \mathrm{L}_2(\mathbb{R}; \mu)$.

It is necessary to distinguish two cases

1. $a_n = 0$ - symmetric case;
2. $a_n \neq 0$ - nonsymmetric case.

In the Hilbert space $\mathcal{H}$ we define the ladder operators $a^{\pm}$ and the number operator N by the formulas

$$\begin{aligned} a^+\Psi_n(x) &= \sqrt{2}b_n\Psi_{n+1}(x), \\ a^-\Psi_n(x) &= \sqrt{2}b_{n-1}\Psi_{n-1}(x), \\ N\Psi_n(x) &= n\Psi_n(x), \quad n \geq 0. \end{aligned}$$

Let $B(N)$ be an operator-valued function such that

$$\begin{aligned} B(N)\Psi_n(x) &= {b_{n-1}}^2\Psi_n(x), \\ B(N+\mathbb{I})\Psi_n(x) &= {b_n}^2\Psi_n(x), \quad n \geq 0. \end{aligned}$$

The next theorem is faithful.

Theorem 1.2. [6] *The operators* $a^{\pm}$*,* N*,* $\mathbb{I}$ *satisfy the following relations*

$$a^-a^+ = 2B(N+\mathbb{I}), \quad a^+a^- = 2B(N), \quad [N, a^{\pm}] = \pm a^{\pm}. \tag{5}$$

Definition. *The associative algebra* $\mathfrak{A}_\Psi$ *generated by the operators* $a^\pm$, N, $\mathbb{I}$ *satisfying the relations of* **theorem 1.2** *and by the commutators of these operators is called the generalized oscillator algebra corresponding to the orthonormal system of polynomials* $\{\Psi_n(x)\}_{n=0}^\infty$ *with recurrence relations* (4).

We will give one useful consequence of the previous theorem. Let $\mathfrak{A}_s$ is the algebra of generalized oscillator corresponding to recurrence relations (4) in the symmetric case ($a_n = 0$) and $\mathfrak{A}_a$ is the algebra of generalized oscillator corresponding recurrence relations (4) in an asymmetric case ($a_n \neq 0$). Then $\mathfrak{A}_s = \mathfrak{A}_a$.

Now we are ready to formulate the problem under consideration. We suppose that there are two families of polynomials $\mathbb{P} = \{P_n(x)\}_{n=0}^\infty$ and $\mathbb{Q} = \{Q_n(x)\}_{n=0}^\infty$ orthogonal with respect to probability measures μ and ν respectively. We suppose that these polynomials satisfy the conditions of **theorem 1.1** and $\beta_n, \widetilde{\beta_n}, \gamma_n, \widetilde{\gamma_n} \in \mathbb{R}$. Let us denote $\mathfrak{A}_P$ and $\mathfrak{A}_Q$ the corresponding algebra of generalized oscillators.

Problem. *We want to describe all pairs of families of orthogonal polynomials* $(\mathbb{P}, \mathbb{Q})$ *for which* $\mathfrak{A}_P = \mathfrak{A}_Q$.

2 Jacobi matrices and the main result

Let $\mathbb{P}$ and $\mathbb{Q}$ are defined in Hilbert spaces $\mathcal{H}_\mu = \mathrm{L}^2(\mathbb{R}; \mu)$ and $\mathcal{H}_\nu = \mathrm{L}^2(\mathbb{R}; \nu)$, respectively. Let these polynomials satisfy the recurrence relations (1) and (3), respectively. In addition, we assume that these polynomials are related to each other by the relation (2). We will also suppose that $\beta_n, \widetilde{\beta_n}, \gamma_n, \widetilde{\gamma_n} \in \mathbb{R}$ and the coefficients β_n, γ_n satisfies the conditions of **theorem 1.1**.

The Jacobi matrices J_P, J_Q corresponding to the RR (1), (3), respectively, can be written in the form

$$J = \begin{bmatrix} A & I_1 \\ I_2 & B \end{bmatrix}, \quad I_1 = \begin{bmatrix} 0 & 0 & 0 & \cdots \\ 0 & 0 & 0 & \cdots \\ 1 & 0 & 0 & \cdots \end{bmatrix}, \quad I_2 = \begin{bmatrix} 0 & 0 & \gamma_3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ \cdots & \cdots & \cdots \end{bmatrix}$$

where the matrix A for sequences $\mathbb{P}$ and $\mathbb{Q}$ have the following form

$$A_P = \begin{bmatrix} \beta_0 & 1 & 0 \\ \gamma_1 & \beta_1 & 1 \\ 0 & \gamma_2 & \beta_2 \end{bmatrix}, \quad A_Q = \begin{bmatrix} \widetilde{\beta}_0 & 1 & 0 \\ \widehat{\gamma}_1 & \widetilde{\beta}_1 & 1 \\ 0 & \widetilde{\gamma}_2 & \widetilde{\beta}_2 \end{bmatrix},$$

and the matrix B equals to

$$B = \begin{bmatrix} \beta_3 & 1 & 0 & 0 & 0 & 0 & \cdots \\ \gamma_2 & \beta_2 & 1 & 0 & 0 & 0 & \cdots \\ 0 & \gamma_3 & \beta_3 & 1 & 0 & 0 & \cdots \\ 0 & 0 & \gamma_2 & \beta_2 & 1 & 0 & \cdots \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \end{bmatrix}.$$

Let us note that elements $(\beta_0, \beta_1, \beta_2, \beta_3)$ and $(\gamma_1, \gamma_2, \gamma_3)$ of the matrix J_P are given, while the elements $(\widetilde{\beta}_0, \widetilde{\beta}_1, \widetilde{\beta}_2)$ as well as coefficients a_1, a_2 in the equation (2) should be defined.

We will consider all 4 cases of the **theorem 1.1**. General relations valid for all four cases have the following form:

$$\begin{aligned} \widetilde{\gamma}_n = \gamma_n,\ n \geq 1; \qquad & \widetilde{\beta}_n = \beta_n,\ n \geq 3; \\ \beta_{2n+1} = \beta_3,\ n \geq 1; \qquad & \gamma_{2n} = \gamma_2,\ n \geq 1; \\ \beta_{2n} = \beta_2,\ n \geq 1; \qquad & \gamma_{2n+1} = \gamma_3,\ n \geq 1; \\ \gamma_n \neq 0; \qquad & a_2 \neq 0. \end{aligned} \tag{6}$$

We use the following notation

$$\begin{gathered} s_1 = \frac{\gamma_3 - \gamma_1 - (\beta_2 - \beta_1)(\beta_3 - \beta_1)}{\gamma_3}, \\ s_2 = \frac{\gamma_2}{\beta_3 - \beta_1}, \quad s_3 = \frac{\gamma_3}{\beta_3 - \beta_1}, \\ w = \frac{a_1}{4s_3} - \frac{\gamma_2}{\gamma_3}, \quad w_\lambda = \frac{\lambda}{(1+\lambda)^2}\,\frac{a_1}{s_3} - \frac{\gamma_2}{\gamma_3}. \end{gathered} \tag{7}$$

We now formulate our main result in terms of Jacobi matrices. Namely, we can prove that all pairs of orthogonal polynomial systems $\{P_n(x)\}_{n=0}^{\infty}$ and $\{Q_n(x)\}_{n=0}^{\infty}$ connected by the linear relation (2), which generate the same algebra of generalized oscillator, can be divided into following eight groups:

The case I $a_1 = 0,\ \beta_1 \neq \beta_3$;
In this case the matrix A_Q and the coefficient a_2 are defined uniquely by the relations

$$\begin{gathered} \widetilde{\beta}_0 = \beta_0 + \frac{(\beta_3 - \beta_1 - \beta_0)\gamma_1}{\gamma_2\gamma_3}\,a_2, \quad a_2 = -s_1 s_3^{\,2}, \\ \widetilde{\beta}_1 = \beta_1 + \frac{a_2}{s_3} - \widetilde{\beta}_0, \quad \widetilde{\beta}_2 = -\frac{a_2}{s_3}, \\ \widetilde{\gamma}_n = \gamma_n,\ n \geq 1, \quad \widetilde{\beta}_n = \beta_n,\ n \geq 3. \end{gathered} \tag{8}$$

The case II $a_1 = 0,\ \beta_1 = \beta_3,\ \gamma_3 = \gamma_1,\ \beta_2 \neq \beta_0$.
In this case the matrix A_Q and the coefficient a_2 are defined uniquely by the relations

$$\begin{gathered} \widetilde{\beta}_0 = \beta_1 - \frac{\gamma_2}{\beta_2 - \beta_0}, \quad \widetilde{\beta}_1 = \beta_0 + \frac{\gamma_2}{\beta_2 - \beta_0}, \\ \widetilde{\beta}_2 = \beta_2,\ a_2 = \gamma_2\,\frac{\beta_1 - \beta_0}{\beta_2 - \beta_0} - \frac{\gamma_2^{\,2}}{(\beta_2 - \beta_0)^2}, \\ \widetilde{\gamma}_n = \gamma_n,\ n \geq 1, \quad \widetilde{\beta}_n = \beta_n,\ n \geq 3. \end{gathered} \tag{9}$$

The case III $a_1 \neq 0,\ \beta_1 \neq \beta_3,\ a_2 = \frac{1}{4}{a_1}^2$.
We denote by w the solution of the equation

$$16{s_3}^2 w^4 + 32 s_2 s_3 w^3 + (16{s_2}^2 + 4{s_3}^2) w^2 + (4s_2 + s_1 s_3) w + s_1 s_2 + \frac{\gamma_2}{\gamma_3}(\beta_2 - \beta_1) = 0$$

such that $a_1 = (4s_3 w + 4s_2) \in \mathbb{R}$ and introduce the quantity

$$C_{\beta,\gamma} = a_2 \left[-\frac{\gamma_1}{\gamma_3} (\beta_2 - a_1(w+1)) + \beta_0(4w^3 + 4w + 1) \right] - a_1 w \left[\beta_0(\beta_1 + \beta_2) + \gamma_1 \right]. \tag{10}$$

In this case, for given w, the matrix A_Q and the coefficients a_1, a_2 are defined uniquely by the relations

$$\widetilde{\beta}_0 = \beta_0 - \frac{C_{\beta,\gamma}}{\gamma_2}, \quad \widetilde{\beta}_1 = \beta_1 + \frac{C_{\beta,\gamma}}{\gamma_2} + a_1 w,$$
$$\widetilde{\beta}_2 = \beta_2 - a_1(w+1), \tag{11}$$
$$a_1 = 4s_3 w + 4s_2, \quad a_2 = \frac{1}{4}{a_1}^2. \tag{12}$$

The case IV $a_1 \neq 0,\ \beta_1 = \beta_3,\ a_2 = \frac{1}{4}\,{a_1}^2$.
Let a_1 be a real solution of the equation

$$\frac{{\gamma_2}^2}{{\gamma_3}^2}\,{a_1}^2 - a_1 \left(\frac{\gamma_2}{\gamma_3} + \frac{\gamma_1}{4\gamma_3} - \frac{1}{4} \right) + \frac{\gamma_2}{\gamma_3}(\beta_2 - \beta_1) = 0.$$

In this case, the matrix A_Q is defined uniquely by the relations

$$\widetilde{\beta}_0 = \beta_0 - \frac{D_{\beta,\gamma}}{\gamma_2}, \quad \widetilde{\beta}_1 = \beta_1 + \frac{D_{\beta,\gamma}}{\gamma_2} + a_1 w, \tag{13}$$
$$\widetilde{\beta}_2 = \beta_2 - a_1(w+1),$$

where $w = -\dfrac{\gamma_2}{\gamma_3}$ and $D_{\beta,\gamma}$ is defined by (10) at $w = -\dfrac{\gamma_2}{\gamma_3}$.

The case V $a_1 \neq 0,\ {a_1}^2 > 4a_2,\ a_2 = \dfrac{\lambda}{(1+\lambda)^2}\,{a_1}^2,\ \beta_1 \neq \beta_3,\ \lambda \in (-1, 1)$.
Let w_λ be a solution of the equation

$$\frac{(1+\lambda)^4}{\lambda^2}\,{s_3}^2 {w_\lambda}^4 + 2\frac{(1+\lambda)^2}{\lambda}\,s_2 s_3 {w_\lambda}^3 + \left(\frac{(1+\lambda)^4}{\lambda^2}\,{s_2}^2 + \frac{(1+\lambda)^2}{\lambda}\,s_3 \right) {w_\lambda}^2 + \left(\frac{(1+\lambda)^2}{\lambda}\,s_2 + s_1 s_3 \right) w_\lambda + s_1 s_2 + \frac{\gamma_2}{\gamma_3}(\beta_2 - \beta_1) = 0, \tag{14}$$

such that

$$a_1 = \frac{(1+\lambda)^2}{\lambda}(s_3 w_\lambda + s_2) \in \mathbb{R}.$$

In this case, the matrix A_Q is defined uniquely by the relations

$$\widetilde{\beta}_0 = \beta_0 - \frac{C_\lambda}{\gamma_2}, \quad \widetilde{\beta}_1 = \beta_1 + \frac{C_\lambda}{\gamma_2} + a_1 w_\lambda, \quad \widetilde{\beta}_2 = \beta_2 - a_1(w_\lambda + 1), \tag{15}$$

where

$$C_\lambda = a_2 \left[-\frac{\gamma_1}{\gamma_3}\left(\beta_2 - a_1(w_\lambda + 1)\right) + \beta_0 \left(\frac{(1+\lambda)^2}{\lambda}({w_\lambda}^2 + w_\lambda) + 1 \right) \right] - a_1 w_\lambda \left[\beta_0(\beta_1 + \beta_2) + \gamma_1\right] \tag{16}$$

$$a_1 = \frac{(1+\lambda)^2}{\lambda}(s_3 w_\lambda + s_2), \quad a_2 = \frac{\lambda}{(1+\lambda)^2}{a_1}^2,$$

and λ is a free parameter.

The case VI $a_1 \neq 0,\ {a_1}^2 > 4a_2,\ a_2 = \dfrac{\lambda}{(1+\lambda)^2}\,{a_1}^2,\ \beta_1 = \beta_3,\ \lambda \in (-1, 1).$

Let a_1 be a real solution of the equation

$$\frac{{\gamma_2}^2}{{\gamma_3}^2}{a_1}^2 - a_1 \left[\frac{\gamma_2}{\gamma_3} + \frac{\lambda}{(1+\lambda)^2}\left(\frac{\gamma_1}{\gamma_3} - 1 \right) \right] + \frac{\gamma_2}{\gamma_3}(\beta_2 - \beta_1) = 0,$$

$$a_2 = \frac{\lambda}{(1+\lambda)^2}{a_1}^2.$$

In this case, the matrix A_Q is defined uniquely by the relations

$$\widetilde{\beta}_0 = \beta_0 - \frac{D_\lambda}{\gamma_2}, \quad \widetilde{\beta}_1 = \beta_1 + \frac{D_\lambda}{\gamma_2} + a_1 w_\lambda, \quad \widetilde{\beta}_2 = \beta_2 - a_1(w_\lambda + 1), \tag{17}$$

where $w_\lambda = -\dfrac{\gamma_2}{\gamma_3}$ and D_λ is defined by the relation (16) for C_λ with $w_\lambda = -\dfrac{\gamma_2}{\gamma_3}$.

The cases VII and VIII $a_1 \neq 0,\ {a_1}^2 < 4a_2,\ a_2 = \dfrac{\lambda}{(1+\lambda)^2}\,{a_1}^2, \quad \lambda = e^{i\theta},\ \theta \in (0, \pi).$

The other relations in the case VII are the same as in the case V, and in the case VIII are the same as in the case VI.

3 Possibility of generalization to the case of $k > 2$

We give few comments about construction all possible pairs of polynomial systems connected by the general linear relation

$$Q_n(x) = P_n(x) + a_1 P_{n-1}(x) + \ldots + a_k P_{n-k}, \quad k \geq 2 \tag{18}$$

for which $\mathfrak{A}_P = \mathfrak{A}_Q$.

From results of the paper [5] we have

1)$\widetilde{\gamma}_n = \gamma_n, \quad n \geq 1;$
2)$a_1, a_2, \ldots, a_k \in \mathbb{R},\ a_k \neq 0,\ \gamma_n \neq 0,\ n \geq 1;$
3) $\widetilde{\beta}_n = \beta_n,\ \widetilde{\gamma}_n = \gamma_n + a_1(\beta_{n-1} - \beta_n) \neq 0, \quad n \geq k+1;$
4) $\gamma_n - \gamma_{n-k} = a_1(\beta_{n-1} - \beta_n), \quad n \geq k+2.$

You should consider two alternative cases:

$$A) \quad a_1 = 0; \qquad B) \quad a_1 \neq 0,\ \beta_n = \beta_k,\ n \geq k.$$

In both cases we have $\gamma_n = \gamma_{n-k}$, $n \geq k+2$.

Then we have to consider all possible cases when some of the coefficients $a_1, a_2, \ldots, a_{k-1}$ are not equal to zero and all others equal to zero. We recall that $a_k \neq 0$. The number of such variants is equal to

$$C_{k-1}^0 + C_{k-1}^1 + \cdots + C_{k-1}^{k-1} = 2^{k-1}.$$

4 The generalized oscillator algebra corresponding to the pair $(\mathbb{P}, \mathbb{Q})$

We discuss the generalized oscillator algebra $\mathfrak{A}_{\mathbb{P}} = \mathfrak{A}_{\mathbb{Q}}$ for considered above pair $(\mathbb{P}, \mathbb{Q})$ of orthogonal polynomials systems connected with each other by the linear relation (18). For the construction of the generalized oscillator algebra $\mathfrak{A}_{\mathbb{P}}$ in the Hilbert space $\mathcal{H}_\mu = \mathrm{L}^2(\mathbb{R}; \mu)$ should be symmetrize the recurrence relations (1) (see [6])

$$xP_n(x) = P_{n+1}(x) + \beta_n P_n(x) + \gamma_n P_{n-1}(x),\ P_0 = 1, \tag{19}$$

with $\gamma_0 = 0$.

With this purpose let us proceed from the system of polynomials $\mathbb{P}$ to the system of polynomials $\Phi = \{\varphi_n(x)\}_{n=0}^{\infty}$ by the formulas

$$P_n(x) = \alpha_n \varphi_n(x),\ \alpha_0 = 1,\ \alpha_n = \sqrt{\gamma_1}(\gamma_2 \ldots \gamma_s)^{\frac{m}{2}}(\gamma_{s+1} \ldots \gamma_{k+1})^{\frac{m-1}{2}},$$

where $n = (m-1)k + s$, $m \geq 1$, $s = 2, \ldots k+1$ and $k \geq 1$ is a fixed integer. This is only possible if $\gamma_1 \neq 0, \ldots, \gamma_{k+1} \neq 0$. Then recurrence relations (19) take the form

$$x\varphi_n(x) = b_n \varphi_{n+1}(x) + \beta_n \varphi_n(x) + b_{n-1}\varphi_{n-1}(x),$$

where

$$\varphi_0 = 1, \quad b_{-1} = 0, \quad b_n = \sqrt{\gamma_{n+1}}, \quad n \geq 0,$$

and

$$\gamma_n = \begin{cases} \gamma_1 & \text{if} \quad n = 1, \\ \gamma_s & \text{if} \quad n = km + s, \quad m \geq 0, \quad s = 2, \ldots k+1. \end{cases} \tag{20}$$

We define the ladder operators $a_\Phi^{\pm}$ and the number operator $\mathbb{N}_\Phi$ in $\mathcal{H}_\mu$ by formulas

$$a_\Phi^+ \varphi_n = \sqrt{2\gamma_{n+1}}\varphi_{n+1},$$
$$a_\Phi^- \varphi_n = \sqrt{2\gamma_n}\varphi_{n-1},$$
$$\mathbb{N}_\Phi \varphi_n = n\varphi_n, \quad n \geq 0.$$

Let $B_\Phi(\mathbb{N}_\Phi)$ be an operator-valued function defined by the following equalities

$$B_\Phi(\mathbb{N}_\Phi)\varphi_n = \gamma_n\varphi_n, \quad n \geq 0.$$

$$B_\Phi(\mathbb{N}_\Phi + I_\mu)\varphi_n = \gamma_{n+1}\varphi_n, \quad n \geq 0.$$

Then the generalized oscillator algebra $\mathfrak{A}_\mathbb{P}$ is generated by the operators $a^\pm_\Phi$, $\mathbb{N}_\Phi$, I_μ satisfying the relations

$$\begin{cases} a^-_\Phi a^+_\Phi = 2B_\Phi(\mathbb{N}_\Phi + I_\mu), \quad a^+_\Phi a^-_\Phi = 2B_\Phi(\mathbb{N}_\Phi), \\ [\mathbb{N}_\Phi, a^\pm_\Phi] = \pm a^\pm_\Phi \end{cases} \tag{21}$$

and by the commutators of these operators.

In this case, the quadratic Hamiltonian

$$H_\Phi = a^-_\Phi a^+_\Phi + a^+_\Phi a^-_\Phi$$

is a selfadjoint operator in the Hilbert space $\mathcal{H}_\mu$.

Orthonormal polynomials $\{\varphi_n(x)\}_{n=0}^\infty$ are eigenfunctions of the operator H_Φ. The corresponding eigenvalues are equal to

$$\lambda_0 = 2\gamma_1, \quad \lambda_n = 2(\gamma_n + \gamma_{n+1}), \quad n \geq 1.$$

In conclusion, let us note that since in our case

$$b_n^2 \neq (a_0 + a_2 n)(1 + n),$$

then according to the results of [9], [10]

$$\dim \mathfrak{A}_\mathbb{P} = \dim \mathfrak{A}_\mathbb{Q} = \infty.$$

Note also that it would be interesting to study the relation of the orthogonality measures μ and ν in the spaces $\mathcal{H}_\mu$ and $\mathcal{H}_\nu$ respectively.

Acknowledgment. EVD grateful to RFBR for financial support under the grant 15-01-03148.

References

[1] V.B.Uvarov, *The connection between systems of polynomials orthogonal with respect to different distribution function*, U.S.S.R. Comput. Math.and Math. Phys., **9**, no.6, 25-36 (1969).

[2] K.H. Kwon, J.H. Lee, F. Marcellan, *Orthogonality of linear combinations of two orthogonal sequences*, Journ. Comput. Appl. Math., **137**, 109-122 (2001).

[3] E. Berriochoa, A. Cachafiero, J.M. Garcia-Amor, *A characterization of the four Chebyshev orthogonal families*, Int. J. Math and Math. Sci. 2005:13, 2071-2079 (2005).

[4] Z.S. Grinshpun, *Differential equation for the Bernstein-Szego orthogonal polynomials*, Differ.Equ. **26**, no.5, 545-550 (1990).

[5] M. Alfaro, F. Marcellan, Ana Pena, M.L. Rezola, *When do linear combinations of orthogonal polynomials yield new sequences of orthogonal polynomials?*, J. Comput. Appl. Math, **233**, 1146-1452 (2010).

[6] V.V. Borzov, *Orthogonal polynomials and generalized oscillator algebras*, Integral Transf. and Special Functions, **12**(2), 115-138 (2001).

[7] T.S. Chihara, *An Introduction to Orthogonal Polynomials*, Gordon and Breach, New York, 1978.

[8] V.V. Borzov, E.V. Damaskinsky, *On representations of generalized oscillator for two sequences of linearly related orthogonal polynomials* Proc. Int. Conf. DAYS on DIFFRACTION 2015, pp.30-33 (2015).

[9] G. Honnouvo, K. Thirulogasanthar, *On the dimensions of the oscillator algebras induced by orthogonal polynomials* J. Math. Phys. **55**, 093511 (2014).

[10] V.V. Borzov, E.V. Damaskinsky, *On dimensions of oscillator algebras* Proc. Int. Conf. DAYS on DIFFRACTION 2014, pp.48-52 (2014).